



10 Old Barn Road · Lake Placid, NY 12946
Technical Support: T: 800 548-7853 · F: 518 523-4513
email: techserv@upstatebiotech.com
Sales Department: T: 800 233-3991 · F: 781 890-7738
Licensing Dept.: 800 310-4659
www.upstatebiotech.com

Certificate of Analysis

Goat Anti-Mouse IgG, HRP-conjugate
(goat polyclonal IgG)
Catalog # 12-349
Lot # 21677

Immunogen: Highly purified whole mouse IgG.

Specificity: Recognizes mouse IgG, both heavy and light chains.

Species Cross-reactivity: Mouse.

Formulation: 500µg of goat IgG, conjugated with horseradish peroxidase, in 250µl 0.02M potassium phosphate, pH 7.2, 0.15M NaCl, 0.01% (w/v) gentamicin sulfate, and 5mg/ml BSA, before the addition of glycerol to 50%. Liquid at -20°C.

Storage and Stability: Stable for 1 year at -20°C from date of shipment.

**FOR RESEARCH USE ONLY
NOT FOR USE IN HUMANS**

Quality Control Testing

Immunoblot Analysis: This lot of antibody was tested at 1:1000, 1:2000, and 1:5000 dilutions in conjunction with anti-Src (Catalog # 05-184) and an EGF-stimulated A431 cell lysate. Previous lots were tested in conjunction with anti-phosphotyrosine (Catalog # 05-321).

Immunoblot Protocol

1. Perform SDS-polyacrylamide gel electrophoresis (SDS-PAGE) on a cell lysate sample (cell lysis buffer: 50mM Tris-HCl, pH7.4; 1% NP-40; 0.25% sodium deoxycholate; 150mM NaCl; 1mM EGTA; 1mM PMSF; 1µg/ml each aprotinin, leupeptin, pepstatin; 1mM Na₃VO₄; 1mM NaF) and transfer the proteins to nitrocellulose. Wash the blotted nitrocellulose twice with water.
2. Block the blotted nitrocellulose in freshly prepared PBS containing 3% nonfat dry milk (PBS-MLK) for 60 minutes at 20-25°C with constant agitation.
3. Incubate the nitrocellulose with **mouse primary antibody**, diluted in freshly prepared PBS-MLK overnight with agitation at 4°C.
4. Wash the nitrocellulose twice with water.
5. Incubate the nitrocellulose in **1:1000-1:5000 Goat Anti-Mouse IgG, HRP-conjugate** in PBS-MLK for 1.5 hours at room temperature with agitation.
6. Wash the nitrocellulose with 4-5 changes of water.
7. Wash the nitrocellulose in PBS-0.05% Tween 20 for 3-5 minutes.
8. Rinse the nitrocellulose in 4-5 changes of water.
9. Use detection method of choice (enhanced chemiluminescence was used).