



28820 Single Oak Drive • Temecula, CA 92590  
Technical Support: T: 800 437-7500 • F: 800 437-7502  
email: techserv@upstate.com  
www.upstate.com

---

## Positive Antigen Controls

### **Non-Stimulated A431 Cell Lysate** (human carcinoma cell line) Catalog # 12-301

**Product Description:** Cellular protein preparation. Cells were lysed in modified RIPA buffer (50 mM Tris-HCl, pH 7.4, 1% NP40, 0.25% sodium deoxycholate, 150 mM NaCl, 1 mM EDTA, 1 mM PMSF, 1 µg/mL aprotinin, 1µg/ml leupeptin, 1µg/ml pepstatin, 1mM Na<sub>3</sub>VO<sub>4</sub>, 1mM NaF) and **diluted with non-reducing sample buffer** (31 mM Tris-HCl, pH 6.8, 5% glycerol, 1% SDS, 0.002% bromphenol blue).

**Use:** Add 2.5 µL of 2-mercaptoethanol/100µl of lysate and boil for 5 minutes to reduce the preparation. Load 20 µg of reduced lysate per lane for immunoblot analysis. This preparation may be used as a positive control for some of Upstate's antibodies.

**Quantity and Formulation:** 100 µg in 100µL of RIPA diluted with non-reducing sample buffer. Concentration: 1 mg/mL. Frozen solution.

**Storage and Stability:** Stable for 1 year at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing.

**FOR RESEARCH USE ONLY  
NOT FOR USE IN HUMANS**