

Certificate of Analysis

28820 Single Oak Drive • Temecula, CA 92590
Technical Support: T: 800 437-7500 • F: 800 437-7502
www.millipore.com

Anti-Fas (human, neutralizing), clone ZB4
(mouse monoclonal IgG₁)
Catalog # 05-338
Lot # DAM1454439

Immunogen: Recombinant human Fas. Clone ZB4.

Specificity: This antibody recognizes the human cell surface antigen Fas expressed in various human cells, including myeloid cells, T lympho-blastoid cells and diploid fibroblasts. It does not induce apoptosis in cell culture.

Species Cross-reactivity: This antibody does not recognize TNF and mouse Fas.

Formulation: 100 µg of ammonium sulfate and protein A purified mouse IgG₁, in 100 µL PBS containing 50% glycerol. Store at -20°C.

Storage and Stability: Stable for 2 years at -20°C from date of shipment.

Handling Recommendations: Upon receipt, and prior to removing the cap, centrifuge the vial and gently mix the solution. Aliquot into microcentrifuge tubes and store at -20°C. **Avoid repeated freeze/thaw cycles, which may damage IgG and affect product performance.** Note: Variability in freezer temperatures below -20°C may cause glycerol-containing solutions to become frozen during storage.

**FOR IN VITRO RESEARCH USE ONLY
NOT FOR USE IN HUMANS OR IN ANIMALS**

Quality Control Testing

Neutralization: 10-500 ng/mL of this lot was incubated for one hour with human Fas-transfected cells and inhibited the anti-Fas (human, activating), clone CH11 (Catalog # 05-201). Cell viability was assessed using a WST-1 assay

Flow Cytometry: Use 5-20 µg/mL.

Immunoblotting: Use 5 µg/mL.

Tested by an outside laboratory

References:

1. Zhang, B., *et al* (2003). Elucidation of molecular events leading to neutrophil apoptosis following phagocytosis: cross-talk between caspase 8, reactive oxygen species, and MAPK/ERK activation. *J Biol Chem* **278**: 28443-54.
2. Lacana, E. and D'Adamio, L. (1999). Regulation of Fas ligand expression and cell death by apoptosis-linked gene 4. *Nat Med* **5**: 542-7.
3. Boirivant, M., *et al* (1996). Stimulated human lamina propria T cells manifest enhanced Fas-mediated apoptosis. *J Clin Invest* **98**: 2616-22.
4. Yonehara, S., *et al* (1994). Involvement of apoptosis antigen Fas in clonal deletion of human thymocytes. *Int Immunol* **6**: 1849-56.