



48 Barn Road • Lake Placid, NY 12946  
Technical Support: T: 800 548-7853 • F: 518 523-4513  
email: techserv@upstate.com  
Sales Department: T: 800 233-3991 • F: 781 890-7738  
Licensing Dept.: 800 310-4659  
www.upstate.com

---

## Positive Antigen Controls

### 3T3/A31 Cell Lysate

(murine fibroblast cell line)

Catalog # 12-305

Lot # 32680

**Product Description:** Cellular protein preparation. Cells were lysed in modified RIPA buffer (50mM Tris-HCl, pH 7.4, 1% NP40, 0.25% sodium deoxycholate, 150mM NaCl, 1mM EDTA, 1mM PMSF, 1µg/ml aprotinin, 1µg/ml leupeptin, 1µg/ml pepstatin, 1mM Na<sub>3</sub>VO<sub>4</sub>, 1mM NaF) and **diluted with non-reducing sample buffer** (31mM Tris-HCl, pH 6.8, 5% glycerol, 1% SDS, 0.002% bromphenol blue).

**Use:** Add 2.5µl of 2-mercaptoethanol/100µl of lysate and boil for 5 minutes to reduce the preparation. Load 20µg of reduced lysate per lane for immunoblot analysis. This preparation may be used as a positive control for some of Upstate's antibodies.

**Quantity and Formulation:** 100µg in 100µl of RIPA diluted with non-reducing sample buffer. Concentration: 1mg/ml. Frozen solution.

**Storage and Stability:** Stable for 6 months at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing.

**FOR RESEARCH USE ONLY  
NOT FOR USE IN HUMANS**