

Certificate of Analysis

Control Histones (+/- Colcemid) (HeLa cell acid extract)

Catalog # 17-306
 Lot # DAM1604869

Kit Components

Control Histones, untreated, (HeLa cell acid extract), Catalog # 13-112. **500µg** packaged in 10 vials, each vial containing **50µg** of precipitated core histones, lyophilized from sterile, distilled water. Lyophilized powder.

Control Histones, colcemid-treated, (HeLa cell acid extract), Catalog # 13-114. **500µg** packaged in 10 vials, each vial containing **50µg** of precipitated core histones, lyophilized from sterile, distilled water. Lyophilized powder.

**FOR RESEARCH USE ONLY
 NOT FOR USE IN HUMANS**

Kit Description

Product Description: Core histones, including histone H1, purified by acid extraction precipitation from log phase untreated and colcemid-treated HeLa cells.


Quantity: 50 assays per kit.

Storage and Stability: Lyophilized: Stable for 2 years at -20°C from date of shipment, when stored with provided desiccant. Reconstituted: Stable for 6 months at -70°C.

Rehydration: Aseptically reconstitute to 1mg/ml with sterile water. Control Histones, untreated may be rehydrated in 2X RSB if used exclusively for immunoblotting. Aliquot to avoid repeated freezing and thawing.

Quality Control Testing

Immunoblot Analysis: Use 5-20µg per lane. 10µg of histones from untreated and colcemid-treated HeLa cells were used as a positive control for immunoblot analysis using 0.5µg/ml anti-phospho-Histone H3 (Ser10) (Catalog # 06-570) and 1:1000 dilution of anti-phospho-Histone H3 (Ser28) (Catalog # 07-145). 1µg/ml anti-phospho-Histone H1 (Catalog # 06-597) was used with a previous lot.



Immunoblot Analysis

Representative lot data. 10µg Control Histones, colcemid-treated (lane 1) or untreated (lane 2), were resolved by electrophoresis, transferred to PVDF and probed anti-phospho-Histone H3 (Ser10) (Catalog # 05-806). Proteins were visualized using a goat anti-mouse (12-349) secondary antibody conjugated to HRP and a chemiluminescence detection system. Arrow indicates phosphorylated Histone H3 (~17kDa).

Immunoblot Protocol

1. Perform SDS-polyacrylamide gel electrophoresis (SDS-PAGE) on **10 μ g Control Histones** and transfer the proteins to PVDF. Wash the blotted PVDF twice with water.
2. Block the blotted PVDF in freshly prepared TBS containing 5% nonfat dry milk (Catalog # 20-200) and 0.05% Tween-20 (TBST-MLK) for 30-60 minutes at room temperature with constant agitation.
3. Incubate the PVDF with an appropriate Histone-specific antibody, diluted in freshly prepared TBST-MLK overnight with agitation at 4°C.
4. Wash the PVDF twice with water.
5. Incubate the PVDF with the appropriate secondary reagent in TBST-MLK for 1.5 hours at room temperature with agitation.
6. Wash the PVDF with water twice.
7. Wash the PVDF in TBS-0.05% Tween 20 for 3-5 minutes.
8. Rinse the PVDF in 4-5 changes of water.
9. Use detection method of choice (enhanced chemiluminescence was used).