

Anti-Phosphotyrosine, clone 4G10

(mouse monoclonal IgG_{2bκ})

Catalog # 05-321

Lot # 19721

Immunogen: Phosphotyramine-KLH.

Antibody Class: IgG_{2bκ} mouse monoclonal antibody produced *in vitro* by mouse-mouse hybridoma 4G10 (FOX-NY [NS-1 derivative] myeloma x spleen cells). Purified by Protein A-Sepharose chromatography.

Formulation: 100mg of mouse monoclonal IgG_{2bκ} in 100ml of 0.01M Tris-HCl, pH 8.0, 0.15M NaCl, 0.02% sodium azide. Protein was determined by a Bradford microtiter protein assay. Liquid.

Storage and Stability: Stable for 6 months at 4°C from date of shipment. **NOTE: DO NOT FREEZE.** For maximum recovery of the product, centrifuge the original vial prior to removing the cap. If the product has accidentally been frozen and thawed, spin it at 13,000 x g for 10 minutes at 4°C. Save the supernatant for application.


**FOR IN VITRO RESEARCH USE ONLY
NOT FOR USE IN HUMANS OR ANIMALS**

Quality Control Testing

Immunoblot Analysis: 0.5-2µg/ml of this lot detected tyrosine-phosphorylated proteins in a modified RIPA lysate from EGF-treated human A431 carcinoma cells.^{1,2,3.}

Included Positive Antigen Control: Catalog #12-302, EGF-stimulated A431 cell lysate is provided as a free positive antigen control for western immunoblotting with this antibody. Aliquot as desired, refreeze immediately, and store at -20°C. The lysate is stable for 6 months at -20°C.

Immunoprecipitation: 2-4µg of this lot can immunoprecipitate quantitatively the phosphotyrosine-containing proteins in the lysate of a confluent culture (10cm dish) of cells expressing an activated tyrosine kinase. To preserve phosphotyrosine, add 0.2mM sodium orthovanadate to the lysis buffer.



Immunoblot Analysis

EGF-stimulated A431 cell lysate was resolved by electrophoresis, transferred to nitrocellulose and probed with anti-phosphotyrosine (1µg/ml). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and a chemiluminescence detection system.

References:

1. Cohen, B., *et al.*, Proc. Natl. Acad. Sci. USA. **87:** 4458-4462, 1990.
2. Druker, B.J., *et al.*, New Eng. J. Med. **321:** 1383-1391, 1989.
3. Kanakura, Y., *et al.*, J. Biol. Chem. **266:** 490-495, 1991.

Immunoprecipitation Protocol

1. Before beginning the immunoprecipitation, dilute the cell lysate to roughly $1\mu\text{g}/\mu\text{l}$ total cell protein in a microcentrifuge tube with PBS.
2. Add **2-4mg of anti-Phosphotyrosine** to $500\mu\text{g}$ - 1mg cell lysate.
3. Gently rock the reaction mixture at 4°C overnight.
4. Capture the immunocomplex by adding $100\mu\text{l}$ of washed Protein A agarose bead slurry ($50\mu\text{l}$ packed beads).
5. Gently rock the reaction mixture at 4°C for 2 hours.
6. Collect the agarose beads by pulsing (5 seconds in the microcentrifuge at $14,000 \times g$), and drain off the supernatant. Wash the beads 3 times with either ice-cold cell lysis buffer or PBS.
7. Resuspend the agarose beads in $50\mu\text{l}$ 2X Laemmli sample buffer.
8. The agarose beads can either be frozen for later use or boiled for 5 minutes. The beads are collected by a microcentrifuge pulse and SDS-PAGE and subsequent immunoblot analysis can be performed on a sample of the supernatant.

Immunoblot Protocol

1. Perform SDS-polyacrylamide gel electrophoresis (SDS-PAGE) on a cell lysate sample (cell lysis buffer: 50mM Tris-HCl, pH 7.4; 1% NP-40; 0.25% sodium deoxycholate; 150mM NaCl; 1mM EDTA; 1mM PMSF; $1\mu\text{g}/\text{ml}$ aprotinin, leupeptin, pepstatin; 1mM Na_3VO_4 ; 1mM NaF) and transfer the proteins to nitrocellulose. Wash the blotted nitrocellulose twice with water.
2. Block the blotted nitrocellulose in freshly prepared PBS containing 3% nonfat dry milk (PBS-MLK) for 20 minutes at 20 - 25°C with constant agitation.
3. Incubate the nitrocellulose with **0.5 - $2\text{mg}/\text{ml}$ of anti-Phosphotyrosine**, diluted in freshly prepared PBS-MLK overnight with agitation at 4°C .
4. Wash the nitrocellulose twice with water.
5. Incubate the nitrocellulose in the secondary reagent of choice (a goat anti-mouse IgG linked to horseradish peroxidase, 1:4000 dilution, was used) in PBS-MLK for 1.5 hours at room temperature with agitation.
6. Wash the nitrocellulose with water twice.
7. Wash the nitrocellulose in PBS-0.05% Tween 20 for 3-5 minutes.
8. Rinse the nitrocellulose in 4-5 changes of water.
9. Use detection method of choice (enhanced chemiluminescence with a 30 second exposure was used).