



Positive Antigen Controls

10 Old Barn Road • Lake Placid, NY 12946
Technical Support: T: 800 548-7853 • F: 518 523-4513
email: techserv@upstate.com
Sales Department: T: 800 233-3991 • F: 781 890-7738
Licensing Dept.: 800 310-4659
www.upstate.com

Non-Stimulated A431 Cell Lysate

(human carcinoma cell line)

Catalog # 12-301

Lot # 28062

Product Description: Cellular protein preparation. Cells were lysed in modified RIPA buffer (50mM Tris-HCl, pH 7.4, 1% NP40, 0.25% sodium deoxycholate, 150mM NaCl, 1mM EDTA, 1mM PMSF, 1µg/ml aprotinin, 1µg/ml leupeptin, 1µg/ml pepstatin, 1mM Na₃VO₄, 1mM NaF) and **diluted with non-reducing sample buffer** (31mM Tris-HCl, pH 6.8, 5% glycerol, 1% SDS, 0.002% bromphenol blue).

Use: Add 2.5µl of 2-mercaptoethanol/100µl of lysate and boil for 5 minutes to reduce the preparation. Load 20µg of reduced lysate per lane for immunoblot analysis. This preparation may be used as a positive control for some of Upstate's antibodies.

Quantity and Formulation: 100µg in 100µl of RIPA diluted with non-reducing sample buffer. Concentration: 1mg/ml. Frozen solution.

Storage and Stability: Stable for 1 year at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing.

**FOR RESEARCH USE ONLY
NOT FOR USE IN HUMANS**