

## **Anti-Human Fas (neutralizes)**

(mouse monoclonal IgG<sub>1</sub>, clone ZB4)

Catalog # 05-338

Lot # 19512

**Immunogen:** Recombinant human Fas. Clone ZB4.

**Specificity:** This antibody recognizes the human cell surface antigen Fas expressed in various human cells, including myeloid cells, T lympho-blastoid cells and diploid fibroblasts. It does not induce apoptosis in cell culture.

**Species Cross-reactivity:** This antibody does not recognize TNF and mouse Fas.

**Formulation: 100mg** of ammonium sulfate and protein A purified mouse IgG<sub>1</sub>, lyophilized from 100µl PBS containing 10mg/ml sucrose. Lyophilized powder. Reconstitute in 100µl distilled water.

**Storage and Stability:** Stable for 2 years at -20°C from date of shipment. Aliquot reconstituted solution to avoid repeated freezing and thawing.

**FOR IN VITRO RESEARCH USE ONLY  
NOT FOR USE IN HUMANS OR IN ANIMALS**

---

### **Quality Control Testing**

**Neutralization:** This lot of antibody, at 250ng/ml or greater, neutralized over 80% of the apoptosis induced by anti-Human Fas, clone CH-11 (Catalog # 05-201, 50ng/ml) treatment of human Jurkat cells.

### Apoptosis Neutralization Assay Protocol

#### A. Day One – Setting up plates and treating cells

1. Add 50µl of cell growth medium (RPMI, 2% FBS) to the wells of a 96 well microtiter plate.
2. Add **50µl of anti-Human Fas, clone ZB4** (at 4x final concentration) in duplicate to row 1.
3. Titrate 50µl from row 1 serially across the plate.
4. Add 50µl of Jurkat cells to each well at a density of  $10^5$  cells/well.
5. Incubate the plate at 37°C, 5% CO<sub>2</sub> for 1 hour to allow antibody/cell interaction.
6. Add 100µl of apoptosis inducing anti-Human Fas, clone CH-11 (Catalog # 05-201), diluted in cell growth medium to 50ng/ml final concentration.
7. Incubate for 20-24 hours at 37°C, 5% CO<sub>2</sub>.

#### B. Day Two - Cell Viability Check - MTT

1. Add 10µl of 5mg/ml MTT in PBS to all wells.
2. Incubate for 2 hours at 37°C, 5% CO<sub>2</sub>.
3. Spin the plate at 2,500 rpm for 5 minutes to pellet the cells.
4. Gently aspirate off the culture media.
5. Add 100µl of acidic isopropanol (0.1N HCl) to each well.
6. Mix until the MTT crystals dissolve, approximately 5 minutes at room temperature.
7. Fill 4 or more wells with 100µl of acidic isopropanol and use to zero the microtiter plate reader
8. Read at 570nm.
9. Compare absorbance of cells that were allowed to grow maximally to cells that were induced to undergo apoptosis in the presence or absence of the neutralizing antibody anti-Human Fas, clone ZB4.