

Anti-Galactocerebroside, clone mGalC

Monoclonal Antibody

Cat. # MAB342

Lot # LV1634994

pack size: 50 µg

Store at -20°C

FOR RESEARCH USE ONLY



Certificate of Analysis

page 1 of 2

Applications	Species Cross-Reactivity	Antibody Isotype	Epitope/Region	Host Species	Molecular Weight	Accession #
WB, ELISA, IH, IC	B, H, M, R, Rb	IgG3	N/A	M	~75 kDa	NM_000153.2

Background

Galactocerebroside (GalC) is a major galactosphingolipid of myelin which plays a role in myelination. GalC is a very useful, specific marker for oligodendroglial lineage.

Presentation

Purified mouse monoclonal in buffer containing 10 mM Potassium Phosphate, 150 mM NaCl, pH 7.4 containing 0.09% sodium azide.

Concentration

1 mg/mL

Specificity

Galactocerebroside (GalC), sulfatide, psychosine and other galactolipids. Cross-reacts with the sulfatide ester of GalC, but to a 16-fold lesser extent. No cross-reactivity with sphingosine, ceramide, mixed ganglioside or glucocerebroside. Binds specifically with oligodendrocytes and Schwann cells.

Immunogen

Synaptic plasma membranes from bovine hippocampus

Method of Purification

Protein A Purified

Storage and Handling

Stable for 6 months at -20°C in undiluted aliquots from date of receipt.

Handling Recommendations: Upon receipt, and prior to removing the cap, centrifuge the vial and gently mix the solution. Aliquot into microcentrifuge tubes and store at -20°C. Avoid repeated freeze/thaw cycles, which may damage IgG and affect product performance..


Control

Neonatal cortex

Quality Control Testing

Evaluated by Western Blot on PC12 lysates.

Western Blotting Analysis: 1:500 dilution of this antibody detected Galactocerebroside on 10 µg of PC12 lysates.



References

1. Chu, Mien-Sheng, *et al.* (2006). *Cell Signal*. 18:519-30.
2. Morishita, H, *et al.* (2004). *Eur J Neurosci*. 20:2843-2847.
3. Sonntag, K., *et al.* (2004). *European Journal of Neuroscience*. 19:1141-1152.
4. Patte-Mensah, C., *et al.* (2003). *J. Neurochemistry*. (2003) 86:1233-1246.


Additional Research Applications

Immunohistochemistry: 0.5-10 µg/mL of a previous lot was used on formalin fixed frozen sections of a previous lot. Can not be used on paraffin embedded tissue sections since the antigen is denatured during embedding and paraffin removal.

Immunocytochemistry: 0.5-10 µg/mL of a previous lot was used on 4% paraformaldehyde, acetic acid or ethanol fixed cultured cells.

ELISA: A previous lot was used on purified galactocerebroside.

Optimal working dilutions must be determined by the end user.



APPLICATION LEGEND: WB Western Blotting ELISA Enzyme-linked Immunosorbent Assay IP Immunoprecipitation IC Immunocytochemistry IF Immunofluorescence IH Immunohistochemistry (Tissue)

SPECIES LEGEND: B Bovine H Human M Mouse R Rat Rb Rabbit WR Most Common Vertebrates

Please visit www.millipore.com for additional product information, test data and references.

Submit your published journal article, and earn credit toward future Millipore purchases. Visit www.millipore.com/publicationrewards to learn more!

upstate | CHEMICON | Linco

THE EXPERTISE OF UPSTATE®, CHEMICON® AND LINCO® IS NOW A PART OF MILLIPORE

Rev. A/2008-07-09/MAB342CA/JM

Anti-Galactocerebroside, clone mGalCCat # MAB342
Lot # LV1634994**PROTOCOL****Western Blot**

1. Perform SDS-polyacrylamide gel electrophoresis (SDS-PAGE) on cell lysate and transfer the proteins to a PVDF membrane. Wash the PVDF membrane twice with water.
2. Block the blotted PVDF membrane in freshly prepared 5% BSA or milk with 0.05% Tween[®]-20 for 1 hour at room temperature with constant agitation.
3. Incubate the PVDF with the recommended dilution of Anti-Galactocerebroside, clone mGalC diluted in freshly prepared 5% BSA or milk for 1 hour at room temperature or overnight with agitation at 2-8°C.
4. Wash the PVDF 3 times with TBST.
5. Incubate the PVDF in the secondary reagent of choice (a Donkey anti-Mouse IgG, HRP conjugate, Catalog # AP192P 1:1000 dilution was used) in 5% milk for 1 hour with agitation at room temperature.
6. Wash the PVDF 3-5 times with TBST.
7. Use Spray and Glow Catalog # 17-373 to visualize results. Use as directed.

RELATED PRODUCTS (specific)

cat #	description
AB142	■ Anti-Galactocerebroside
AP124P	■ Goat anti-Mouse IgG, Peroxidase Conjugated, H+L

RELATED PRODUCTS (non-specific)

cat #	description
IPVH00010	■ Immobilon-P 26.5 cm x 3.75 m Roll PVDF 0.45 µm
IPFL00010	■ Immobilon-FL 26.5 cm x 3.75 m Roll PVDF 0.45 µm
IPVH07850	■ Immobilon-P 7 x 8.4 cm PVDF 0.45 mm (sheet) 50/pk
ISEQ00010	■ Immobilon-P SQ 26.5 cm x 3.75 m 1 roll PVDF 0.2 µm
ISEQ07850	■ Immobilon-P 7 x 8.4 cm PVDF 0.2 mm (sheet) 50/pk
IPFL07810	■ Immobilon-FL 7 x 8.4 cm PVDF 0.45 mm (sheet) 10/pk
WBKLS0100	■ Immobilon Western Chemilum HRP Substrate 100 mL
17-373	■ Spray & Glow™ ECL WB Detection System 1 ea
2060	■ Re-Blot Western Blot Recycling Kit
2500	■ Re-Blot Plus Western Blot Recycling Kit
B2080-175GM	■ Blot Quick Blocker Membrane Blocking Agent 175G

■ antibodies ■ Multiplex products ■ biotools ■ cell culture ■ enzymes ■ kits ■ proteins/peptides ■ siRNA/cDNA products

Please visit www.millipore.com for additional product information, test data and references

28820 Single Oak Drive • Temecula, CA 92590

Technical Support: T: 1-800-MILLIPORE (1-800-645-5476) • F: 1-800-437-7502

FOR RESEARCH USE ONLY. Not for use in diagnostic or therapeutic applications. Not for human or animal consumption. Purchase of this Product does not include any right to resell or transfer, either as a stand-alone product or as a component of another product. Any use of this Product for purposes other than research is strictly prohibited without prior written authorization from an authorized officer of Millipore Corporation.

Upstate®, Chemicon®, Linco® and all other trademarks are owned by Millipore Corporation. Copyright ©2008-2009 Millipore Corporation. All rights reserved.



We Buy 100% Certified Renewable Energy