

---

## Certificate of Analysis

### Protein A Agarose

(10ml packed beads)

Catalog # 16-125

Lot # 22929

**Description and Formulation:** 10ml packed beads containing sufficient covalently-linked Protein A for a binding capacity of  $20 \pm 2$  mg/ml human IgG, as determined by an independent laboratory. Protein A is linked by a proprietary method to minimize leakage to less than 3ng/ml as determined by ELISA. Provided as a 50% gel slurry for a final volume of 20ml. Suspended water containing 0.01% thimerosal.

**Physical Form:** Liquid suspension. Prior to use, wash the agarose beads with an appropriate buffer to remove the thimerosal.

**Storage and Stability:** Stable for 1 year at 4°C from date of shipment.

**FOR RESEARCH USE ONLY  
NOT FOR USE IN HUMANS**

---

### Quality Control Testing and Research Applications

Immunoprecipitation: Previous lots were tested by using 100 $\mu$ l of the gel slurry with monoclonal anti-Phosphotyrosine (Catalog # 05-321) to immunoprecipitate phosphotyrosine containing proteins from an EGF-stimulated A431 cell lysate (Catalog # 12-302).

Affinity Purification of IgG: A previous lot was tested by using 5ml of the gel slurry to quantitatively capture the IgG from 10ml of rabbit antiserum.

---

### Immunoprecipitation Protocol

1. Prepare a cell lysate at a concentration of about 1 $\mu$ g/ $\mu$ l of protein and add 500 $\mu$ g-1mg to a microfuge tube.
2. Add an appropriate amount of primary antibody to the tube.
3. Gently rock the reaction mixture at 4°C overnight.
4. Capture the immunocomplex by adding 100 $\mu$ l (50 $\mu$ l packed beads) of washed Protein A agarose bead slurry.
5. Gently rock the reaction mixture at 4°C for 2 hours.
6. Collect the agarose beads by pulsing (5 seconds in the microcentrifuge at 14,000 x g), and drain off the supernatant.
7. Wash the beads 3 times with either ice-cold cell lysis buffer or PBS.
8. Resuspend the agarose beads in 60 $\mu$ l 2X Laemmli sample buffer and boil for 5 minutes. Collect the beads by a microcentrifuge pulse. SDS-PAGE and subsequent immunoblot analysis can be performed on a sample of the supernatant, or the agarose beads can then be frozen for later use and reboiled for 5 minutes prior to SDS-PAGE.