



Certificate of Analysis

EZ ChIP™ Chromatin Immunoprecipitation Kit Catalog # 17-371 Lot # DAM1604878

Kit Components

ChIP Blocked Protein G Agarose Catalog #16-201D. One vial containing **1.5 mL** packed beads ChIP Blocked with 1.5 mg BSA and approximately 4.5 mg recombinant Protein G. Provided as a 50% gel slurry for a final volume of **3 mL** per vial. Suspended in TE buffer, pH 8.0, containing 0.05% sodium azide. Liquid suspension.

ChIP Dilution Buffer, Catalog # 20-153. One vial containing **24 mL**.

Low Salt Immune Complex Wash Buffer, Catalog # 20-154. One vial containing **24 mL**.

High Salt Immune Complex Wash Buffer, Catalog # 20-155. One vial containing **24 mL**.

LiCl Immune Complex Wash Buffer, Catalog # 20-156. One vial containing **24 mL**.

TE Buffer, Catalog # 20-157. Two vials, each containing **24 mL**.

0.5M EDTA, Catalog # 20-158. One vial containing **250 µL**.

5M NaCl, Catalog # 20-159. One vial containing **500 µL**.

SDS Lysis Buffer, Catalog # 20-163. One vial containing **10 mL**.

1M Tris-HCl, pH 6.5, Catalog # 20-160. One vial containing **500 µL**.

10X PBS, Catalog # 20-281. One vial containing **24 mL**.

10X Glycine, Catalog # 20-282. One vial containing **11 mL**.

Protease Inhibitor Cocktail II, Catalog # 20-283. Two vials, each containing **100 µL** of 200X Protease Inhibitor Cocktail II in DMSO.

RNase A, Catalog # 20-297. One vial containing **600µg** of RNase A in **60 µL** of sterile water.

Proteinase K, Catalog # 20-298. One vial containing **600 µg** of Proteinase K in **60 µL**.

1M NaHCO₃, Catalog # 20-296. One vial containing **600 µL**.

Control Primers, Catalog # 22-004, One vial containing **75 µL** of 5µM of each control primer specific for human GAPDH.

FOR: 5'-TACTAGCGGTTTTACGGGCG-3'
REV: 5'-TCGAACAGGAGGAGCAGAGAGCGA-3'

Anti-RNA Polymerase II, clone CTD4H8, Catalog # 05-623B. One vial containing **25µg** of Anti-RNA Polymerase II, clone CTD448.

Normal Mouse IgG, Catalog # 12-371B. One vial containing **25µg** of normal mouse IgG.

20% SDS, Catalog # 20-280. One vial containing **241 µL**.

Spin Filters, Catalog # 20-290. One bag containing 22 Spin Filters in Collection Tubes.

Collection Tubes, Catalog # 20-291. One bag containing 22 Collection Tubes.

Bind Reagent A, Catalog # 20-292. One vial containing **25 mL** of Bind Reagent A.

Wash Reagent B, Catalog # 20-293. One vial containing **12.5 mL** of Wash Reagent B.

Elution Reagent C, Catalog # 20-294. One vial containing **1.5 mL** of Elution Reagent C.

**FOR RESEARCH USE ONLY
NOT RECOMMENDED OR INTENDED FOR DIAGNOSIS OF DISEASE IN HUMANS
DO NOT USE IN HUMANS**

Kit Description

Quantity: Two boxes containing the necessary reagents to perform 22 chromatin immunoprecipitation (ChIP) assays. This kit also contains reagents to generate chromatin from ten 10-cm plates to allow for multiple immunoprecipitations from the same chromatin preparation.


Storage and Stability: Upon receipt, store components at the temperatures indicated on the labels. Kit components are stable for 1 year from date of shipment when stored as directed.

Use: The EZ ChIP™ kit contains reagents optimized for immunoprecipitation of chromatin from mammalian cells, including controls to ensure successful performance of this assay. Detection of the DNA region, gene or promoter of interest in immunoprecipitated chromatin must be empirically determined by the researcher. Quantitative PCR using promoter-specific primers is recommended. The EZ ChIP™ kit has all the necessary buffers and reagents to perform successful chromatin immunoprecipitation assays. However, careful attention must be paid to the details of the instructions. Follow all instructions carefully, especially with regard to incubation times and temperatures.

Please refer to the Instruction Manual for further information and a detailed assay protocol.

Quality Control Testing

PCR Analysis of Chromatin Immunoprecipitation:
Representative lot data. Chromatin immunoprecipitation was performed using chromatin from HeLa cells and either anti-RNA Polymerase II (Catalog # 05-623B) or Normal Mouse IgG (Catalog # 12-371B) as the immunoprecipitating antibody. Purified DNA was then analyzed by PCR using Control Primers specific for the GAPDH promoter. PCR product was observed in the anti-RNA Polymerase II ChIP (lane 3) and not in the Normal Mouse IgG ChIP (lane 2). GAPDH promoter specific DNA was also observed in the Input (lane 4) and not in the "No DNA" PCR control (lane 1).



General References:

1. Das, Partha M, *et al* (2004). Chromatin immunoprecipitation assay. *BioTechniques* **37**: 961-9.
2. Luo, R X, *et al* (1998). Rb interacts with histone deacetylase to repress transcription. *Cell* **92**: 463-73.
3. Braunstein, M, *et al* (1996). Efficient transcriptional silencing in *Saccharomyces cerevisiae* requires a heterochromatin histone acetylation pattern. *Mol Cell Biol* **16**: 4349-56..
4. Buck, Michael J and Lieb, Jason D (2004). ChIP-chip: considerations for the design, analysis, and application of genome-wide chromatin immunoprecipitation experiments. *Genomics* **83**: 349-60.
5. Bernstein, Bradley E, *et al* (2004). The use of chromatin immunoprecipitation assays in genome-wide analyses of histone modifications. *Meth Enzymol* **376**: 349-60.
6. Weinmann, Amy S, *et al* (2002). Isolating human transcription factor targets by coupling chromatin immunoprecipitation and CpG island microarray analysis. *Genes Dev* **16**: 235-44.
7. Manabe, I and Owens, G K (2001). CArG elements control smooth muscle subtype-specific expression of smooth muscle myosin in vivo. *J Clin Invest* **107**: 823-34.
8. Cervoni, N and Szyf, M (2001). Demethylase activity is directed by histone acetylation. *J Biol Chem* **276**: 40778-87.

Unless otherwise stated in our catalog or other company documentation accompanying the product(s), our products are intended for research use only and are not to be used for any other purpose, which includes but is not limited to, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses or any type of consumption or application to humans or animals.

©2007: Millipore Corporation. All rights reserved. No part of these works may be reproduced in any form without permission in writing.