

Anti-Caspase 3
(rabbit polyclonal IgG)
Catalog # 06-735
Lot # 19227

Immunogen: Human full-length Caspase 3 fusion protein containing a histidine-6 tag.

Specificity: Recognizes full-length Caspase 3 (Yama/Apopain) at 32kDa and proteolytic fragments as demonstrated after *in vitro* cleavage of recombinant Caspase 3.

Species Cross-reactivity: Human, mouse and rat.

Formulation: 200µg protein A purified IgG in 200µl of 0.07M Tris-glycine, pH 7.4, 0.105M NaCl, 0.035% sodium azide with 30% glycerol. Liquid at -20°C.


Storage and Stability: Stable for 2 years at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial prior to removing the cap.

FOR RESEARCH USE ONLY
NOT FOR USE IN HUMANS

Quality Control Testing

Immunoblot Analysis: 0.5-2µg/ml of this lot detected the 32kDa precursor of Caspase 3 in RIPA lysates from non-stimulated human A431 cells; previous lots detected Caspase 3 in RIPA lysates of mouse 3T3 and rat PC12 cells.

Included Positive Antigen Control: Catalog # 12-301, non-stimulated A431 cell lysate. Use 20µg per lane for minigels.



Immunoblot Analysis
Non-stimulated A431 cell lysate was resolved by electrophoresis, transferred to nitrocellulose and probed with anti-Caspase 3 (1µg/ml). Proteins were visualized using a goat anti-rabbit secondary antibody conjugated to HRP and a chemiluminescence detection system. Arrow indicates the precursor of Caspase 3 (32kDa).

Immunoblot Protocol

1. Perform SDS-polyacrylamide gel electrophoresis (SDS-PAGE) on a cell lysate sample (cell lysis buffer: 50mM Tris-HCl, pH7.4; 1% NP-40; 0.25% sodium deoxycholate; 150mM NaCl; 1mM EGTA; 1mM PMSF; 1µg/ml aprotinin, leupeptin, pepstatin; 1mM Na₃VO₄; 1mM NaF) and transfer the proteins to nitrocellulose. Wash the blotted nitrocellulose twice with water.
2. Block the blotted nitrocellulose in freshly prepared PBS containing 3% nonfat dry milk (MLK) for 60 minutes at 20-25°C with constant agitation.
3. Incubate the nitrocellulose with **0.5-2µg/ml of anti-Caspase 3**, diluted in freshly prepared PBS-MLK overnight with agitation at 4°C.
4. Wash the nitrocellulose twice with water.
5. Incubate the nitrocellulose in the secondary reagent of choice (a **goat anti-rabbit** HRP conjugated IgG, 1:3000 dilution was used) in PBS-MLK for 1.5 hours at room temperature with agitation.
6. Wash the nitrocellulose with water twice.
7. Wash the nitrocellulose in PBS-0.05% Tween 20 for 3-5 minutes.
8. Rinse the nitrocellulose in 4-5 changes of water.
9. Use detection method of choice (enhanced chemiluminescence was used). Detection may require long exposure times (10-30 minutes) in some systems.