

Anti-Myc Tag, clone 4A6

(mouse monoclonal IgG₁)

Catalog # 05-724

Lot # 24566

Immunogen: KLH-conjugated, synthetic peptide corresponding to amino acids 410-420 (MEQKLISEEDL) of human Myc.

Specificity: Recognizes and is specific for recombinant proteins containing the Myc epitope tag (EQKLISEEDL) in a variety of sequence contexts. Also recognizes human Myc.

Species Cross-reactivity: Human. Other species cross-reactivity not tested.

Formulation: 200µg of protein G purified mouse IgG₁ in 414µl of 0.1M Tris-glycine, pH 7.4, 0.15M NaCl, 0.05% sodium azide before the addition of glycerol to 30%. Liquid at -20°C.

Storage and Stability: Stable for 2 years at -20°C from date of shipment. For maximum recovery of product, centrifuge the vial prior to removing the cap.


**FOR IN VITRO RESEARCH USE ONLY
NOT FOR USE IN HUMANS OR ANIMALS**

Quality Control Testing

Immunoblot Analysis: 0.5-2.0µg/ml of this lot detected Myc-tagged recombinant protein in sequence contexts not well recognized by anti-Myc Tag, clone 9E10 (Catalog # 05-419).


Additional Research Applications

Immunoprecipitation: An independent laboratory has reported that this antibody immunoprecipitates Myc-tagged protein from transfected cells.



Immunoblot Analysis

Panel A. Lysates from NIH/3T3 cells transfected with either empty vector (lanes 1, 4), PP2A A subunit containing the Myc epitope tag sequence MEQKLISEEDLLRKRGS (lanes 2, 5), or PP2A A subunit containing the Myc epitope tag sequence MEQKLISEEDLNGST (lanes 3, 6) were resolved by electrophoresis, transferred to nitrocellulose and probed with anti-Myc Tag, clone 4A6 (0.5µg/ml, lanes 1-3) or anti-Myc Tag, clone 9E10 (1µg/ml, Catalog # 05-419, lanes 4-6). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and a chemiluminescence detection system. Arrow indicates Myc-tagged PP2A A subunit (~63kDa).



Panel B. Longer exposure of lanes 4-6 demonstrating weak detection by anti-Myc Tag, clone 9E10 of the Myc-tagged protein containing the epitope tag sequence MEQKLISEEDLNGST

Immunoblot Protocol

1. Perform SDS-polyacrylamide gel electrophoresis (SDS-PAGE) on a transfected cell lysate sample (cell lysis buffer: 50mM Tris-HCl, pH 7.4; 1% NP-40; 0.25% sodium deoxycholate; 150mM NaCl; 1mM EDTA; 1mM PMSF; 1 μ g/ml each aprotinin, leupeptin, pepstatin; 1mM Na₃VO₄, 1mM NaF) and transfer the proteins to nitrocellulose. Wash the blotted nitrocellulose twice with water.
2. Block the blotted nitrocellulose in freshly prepared 3% nonfat dry milk (Catalog # 20-200) in TBS (TBS-MLK) for 1 hour at room temperature with constant agitation.
3. Incubate the nitrocellulose with **0.5-2.0 μ g/ml of anti-Myc Tag, clone 4A6**, diluted in freshly prepared TBS-MLK for 2 hours at room temperature with constant agitation.
4. Wash the nitrocellulose twice with water.
5. Incubate the nitrocellulose in the secondary reagent of choice (a goat anti-mouse HRP conjugated IgG, Catalog # 12-349, 1:5000 dilution was used) in TBS-MLK for 30 minutes at room temperature with agitation.
6. Wash the nitrocellulose twice with water.
7. Wash the nitrocellulose in TBS-0.05% Tween 20 for 3-5 minutes.
8. Rinse the nitrocellulose in 4-5 changes of water.
9. Use detection method of choice (enhanced chemiluminescence was used).