

Anti-Rabbit IgG, HRP-conjugated

(goat polyclonal IgG)

Catalog #12-348

Lot # 17520

Immunogen: Highly purified whole rabbit IgG.

Specificity: Specific for rabbit IgG heavy and light chains.

Species Cross-reactivity: Rabbit

Formulation: 500mg of goat IgG, conjugated with horseradish peroxidase, in 250ml 0.02M potassium

phosphate, 0.15M NaCl, pH 7.2, with 10mg/ml BSA, 50% glycerol and 0.01% (w/v) gentamicin sulfate. Liquid at -20°C.

Storage and Stability: Stable for 1 year at -20°C from date of shipment.

**FOR RESEARCH USE ONLY
NOT FOR USE IN HUMANS**

Quality Control Testing

Immunoblot Analysis: This lot of antibody was tested at 1:1000, 1:3000 and 1:5000 dilutions in conjunction with a rabbit polyclonal anti-MAP Kinase R2 (Catalog # 06-182) antibody to detect MAPK (42-44kDa) in a mouse 3T3 cell lysate.

Immunoblot Protocol

1. Perform SDS-polyacrylamide gel electrophoresis (SDS-PAGE) on a cell lysate sample (cell lysis buffer: 50mM Tris-HCl, pH7.4; 1% NP-40; 0.25% sodium deoxycholate; 150mM NaCl; 1mM EGTA; 1mM PMSF; 1µg/ml each aprotinin, leupeptin, pepstatin; 1mM Na₃VO₄; 1mM NaF) and transfer the proteins to nitrocellulose. Wash the blotted nitrocellulose twice with water.
2. Block the blotted nitrocellulose in freshly prepared PBS containing 3% nonfat dry milk (PBS-MLK) for 90 minutes at 20-25°C with constant agitation.
3. Incubate the nitrocellulose with **the rabbit primary antibody**, diluted in freshly prepared PBS-MLK overnight with agitation at 4°C.
4. Wash the nitrocellulose twice with water.
5. Incubate the nitrocellulose in **1:1000-1:5000 goat anti-rabbit IgG, HRP-conjugated** in PBS-MLK for 1.5 hours at room temperature with agitation.
6. Wash the nitrocellulose with water twice.
7. Wash the nitrocellulose in PBS-0.05% Tween 20 for 3-5 minutes.
8. Rinse the nitrocellulose in 4-5 changes of water.
9. Use detection method of choice (enhanced chemiluminescence was used).