

ESGRO[®] Supplement (10⁶ Units)

Cell Culture Supplement

Cat. # ESG1106

Lot # 7240-102E

FOR RESEARCH USE ONLY

pack size: 10⁶ units

Store at 2-8°C
DO NOT FREEZE



Certificate of Analysis

page 1 of 2

Description

ESGRO medium supplement is an optimized formulation of mouse Leukemia Inhibitory Factor (LIF), which promotes self-renewal and long-term maintenance of pluripotency of mouse embryonic stem (mES) cells by suppressing spontaneous differentiation. Each lot of ESGRO supplement undergoes stringent quality control testing to ensure consistent inhibition of mES cell differentiation.

Purity and Sterility

The active component mLIF has been shown to be >95% pure by SDS-PAGE. ESGRO supplement is supplied 0.22 micron filtered, and tested negative in both aseptic and microplasmic tests. Endotoxin level is less than 0.1 ng per µg of LIF.

Activity

ESGRO supplement is assessed both on mES cells and on murine M1 myeloid leukemic cells. A standard of 50 Units is defined as the concentration of ESGRO supplement in 1.0 mL of tissue culture medium that induces the differentiation of 50% of M1 colonies (Metcalf, 1988).

Embryonic Stem Cell Assay: Differentiation inhibition at 1000 units/mL

Murine myeloid leukemic, M1 Assay: Specific Activity > 10⁸ units/mg

Presentation

ESGRO supplement is supplied in liquid form as 10⁶ Units in 1.0 mL of phosphate buffered saline with 1% w/v bovine serum albumin BSA as a carrier for stability. 1x10⁶ units is sufficient to treat 1L of ES cell culture media.

Storage and Handling

ESGRO supplement is shipped on cold packs.. Refer to the product vial label for exact expiry information. ESGRO supplement is stable in the concentrated form or diluted in sterile tissue culture media, with no loss of activity on ES cells up to expiration date. For long term storage it is recommended that ESGRO concentrate be stored at 2-8°C. Freeze thawing will reduce potency.

It is recommended that prior to use, ESGRO supplement should be diluted in sterile tissue culture media and aliquoted to a convenient concentration, then stored at 2-8°C. Freeze-thawing should be avoided.

ESGRO supplement is stable for a minimum of 7 days at 37°C, 5% CO₂ incubator during the culture of ES cells.

Warranty

The highest standards of quality control are used in the manufacture of this product. No warranty is provided that the sale or use of the product either alone, in combination with other products, or in the operation of any process, will not infringe patent, intellectual property or any other rights of third parties. This product is supplied for research purposes only. It is not for human therapeutic and/or diagnostic use.

NOTE: ESGRO supplement should not be allowed to come in contact with ruminant animals or swine.

Suggested Protocols

ES Cells:

For routing mouse ES cell culture, 1000 U of ESGRO supplement per 1.0 mL of tissue culture media is sufficient to maintain ES cells with a stem cell phenotype. Similar concentrations of mLIF have also been used for germline transmission of genetically altered ES cells (Ziilstra, 1989).

At the recommended concentration 10⁷ units of ESGRO supplement is sufficient for 10.0 L of tissue culture media and 10⁶ units of ESGRO supplement is sufficient for 1.0 L of tissue culture media.

Please visit www.millipore.com for additional product information, test data and references.

Submit your published journal article, and earn credit toward future Millipore purchases.

Visit www.millipore.com/publicationrewards to learn more!

upstate | CHEMICON | *Linco*

THE EXPERTISE OF UPSTATE[®], CHEMICON[®] AND LINCO[®]
IS NOW A PART OF MILLIPORE

Rev. A/2009-8-20/ESG1106CA/AN

References and Suggested Readings:

1. Smith AG., Heath JK., Donaldson DD., et al. "Inhibition of pluripotential embryonic stem cell differentiation by purified polypeptides." Nature 336:688-90, 1988.
2. Williams RL., Hilton DJ., Pease S., et al. "Myeloid leukemia inhibitory factor maintains the developmental potential of embryonic stem cells. Nature, 336:686-7, 1988.
3. Zijlstra M, Li E, Sajjadi F, Subramani S, Jaenisch R. "Germ-line transmission of a disrupted $\beta 2$ -microglobulin gene produced by homologous recombination in embryonic stem cells." Nature 342:435-8, 1989.
4. Metcalf D., Hilton DJ., Nicola NA. "Clonal analysis of the actions of the murine leukaemia inhibitory factor on leukemic and normal murine haemopoietic cells. Leukemia 2:216-21, 1988.

Please Read Before Opening Product

Leukemia Inhibitory Factor ("LIF") is protected by US Patents No. 5,166,065, 5,187,077, 5,427,925, 5,750,654 and 6,261,548 owned by AMRAD Corporation Ltd. and licensed exclusively to Millipore for research uses.

LABEL LICENSE - USE RESTRICTED TO RESEARCH USE ONLY

Leukemia Inhibitory Factor ("LIF") is protected by US Patents No. 5,166,065, 5,187,077, 5,427,925, 5,750,654 and 6,261,548 owned by AMRAD Corporation Ltd. and licensed exclusively to Millipore for research uses. The use of this product is limited for research uses only and not for any commercial purposes, such as making or developing other research products from this product or use of this product as a component in a research kit. This product is not to be used in humans or for any diagnostic purposes.

The purchaser may refuse to accept the conditions of this label license by returning this product unopened and will receive a full refund of the purchase price paid.

For-profit entities or commercial research entities purchasing this product shall be required to execute a separate commercial license with Millipore within three months of purchase. Researchers may not transfer this product or its derivatives to researchers performing commercial research at other Nonprofit Organizations, or to For-Profit Organizations, without the express written consent of Millipore, and without those entities having an express license from Millipore for the use of the product. Other than as specifically provided herein, a transferee of this product shall have no right to transfer this product, or its derivatives to any other person or entity.

RELATED PRODUCTS

cat #		description
ESG1107	■	ESGRO mLIF Medium Supplement, 10 ⁷ units
SLM-220-B	■	ES Cell Qualified DMEM
ES-009-B	■	ES Cell Qualified FBS
TMS-002-C	■	L-Glutamine
ES-007-E	■	2-mercaptoethanol
ES-008-D	■	Nucleosides
TMS-001-C	■	Non-Essential Amino Acids
TMS-AB2-C	■	Penicillin-Streptomycin
LIF2005	■	Recombinant Mouse Leukemia Inhibitory Factor, 5 μ g
LIF2010	■	Recombinant Mouse Leukemia Inhibitory Factor, 10 μ g

■ antibodies ■ Multiplex products ■ biotools ■ cell culture ■ enzymes ■ kits ■ proteins/peptides ■ siRNA/cDNA products

Please visit www.millipore.com for additional product information, test data and references

28820 Single Oak Drive • Temecula, CA 92590

Technical Support: T: 1-800-MILLIPORE (1-800-645-5476) • F: 1-800-437-7502



We Buy 100% Certified Renewable Energy

FOR RESEARCH USE ONLY. Not for use in diagnostic or therapeutic applications. Not for human or animal consumption. Purchase of this Product does not include any right to resell or transfer, either as a stand-alone product or as a component of another product. Any use of this Product for purposes other than research is strictly prohibited without prior written authorization from an authorized officer of Millipore Corporation.

Upstate®, Chemicon®, Linco® and all other trademarks are owned by Millipore Corporation. Copyright ©2008-2009 Millipore Corporation. All rights reserved.