



Fischer Building, Gemini Crescent Dundee,
DD2 1SW, UK
T: +44(0) 1382 561600
F: +44(0) 1382 561601
Technical Support: (US/Canada)
T: 800 437-7500 • F: 800 437-7502
www.millipore.com

Certificate of Analysis

Lck, active

(Recombinant enzyme expressed in Sf21 insect cells)

Catalogue # 14-442

Lot # 1610220

Product Description: N-terminal 6His-tagged, recombinant full-length human Lck expressed in Sf21 insect cells. Purified using Ni²⁺/NTA agarose. Purity 85% by SDS-PAGE and Coomassie blue staining. MW = 59kDa.

Formulation: 10µg of enzyme in 5.2µl of 50mM Tris/HCl, pH7.5, 0.1mM EGTA, 150mM NaCl, 1mM benzamidine, 0.03% Brij-35, 0.2mM PMSF, 0.1% 2-mercaptoethanol, 5% glycerol. Frozen solution.

Storage and Stability: Stable for 1 year at -70°C from date of shipment. For maximum recovery of product, centrifuge original vial prior to removing the cap.

Specific Activity (lot# 1610220): 375U/mg, where one unit of Lck activity is defined as 1nmol phosphate incorporated into 250µM cdc 2 peptide per minute at 30°C with a final ATP concentration of 100µM.

Handling Recommendations: Rapidly thaw the vial under cold water and immediately place on ice. Aliquot unused material into pre-chilled microcentrifuge tubes and immediately snap-freeze the vials in liquid nitrogen prior to re-storage at -70°C.



**FOR IN VITRO RESEARCH USE ONLY
NOT FOR USE IN HUMANS OR ANIMALS**

Quality Control Testing

Kinase Assay: 192–480ng of this lot of enzyme phosphorylated 250µM cdc2 peptide in the assay described on page two. Assay background was subtracted from the actual counts to yield the results shown below.

MS Tryptic Fingerprint: Confirmed identity as Lck with 61% amino acid coverage of the translated sequence listed on page three.

SDS-PAGE and Coomassie Stain: Representative gel from this lot. Purity was assessed by SDS-PAGE and Coomassie blue staining using 3µg of active Lck.



Kinase Assay Protocol

Stock Solutions:

1. **4 x Reaction Buffer:** 200mM Tris/HCl pH7.5, 0.4mM EGTA, 0.4mM Na₃VO₄.
2. **cdc2 Substrate Peptide:** Use a final concentration of 250μM. Make a 2.5mM stock. Add 2.5μl of stock per assay point.
3. **Lck, active:** Dilute with 20mM MOPS/NaOH pH7.0, 1mM EDTA, 5% glycerol, 0.01% Brij-35, 0.1% 2-mercaptoethanol, 1mg/ml BSA. Use 192–480ng per assay point.
4. **[γ-³³P]ATP:** 2.5 x magnesium acetate/[γ-³³P]ATP cocktail: 25mM MgAc and 0.25mM ATP to which is added [γ-³³P]ATP (specific activity approximately 500 - 800cpm/pmol as required.)

Assay Procedure (96 well plate format):

1. Add 6.25μl of 4 x reaction buffer per assay to wells.
2. Add 2.5μl of substrate peptide **cdc2**.
3. Add **2.5μl (192–480ng) Lck, active**.
4. Add 3.75μl of dH₂O.
5. Add 10μl of diluted [γ-³³P]ATP mixture.
6. Incubate for 10 minutes at 30°C.
7. Stop the reaction by adding 5μl of 3% phosphoric acid.
8. Transfer a 10μl aliquot onto the appropriate area of a **P30 Filtermat**.
9. Wash the filtermat twice for 5 minutes with 75mM phosphoric acid.
10. Wash the filtermat once for 2 minutes with methanol.
11. Transfer the filtermat to a sealable plastic bag and add 4ml of scintillation cocktail.
12. Read in a scintillation counter. Compare cpm of enzyme samples with cpm of control samples that contain all assay components plus 1μl of 30% phosphoric acid (background cpm.)

Lck Sequence Information

Protein	Human Lck
Tags	N-terminal 6His
Native sequence	G9 of the recombinant protein is equivalent to G1 of human Lck
Accession number	Swiss Prot P06239

Recombinant Lck amino acid sequence:

```
1 MHHHHHMGCGCSSHPEDDW MENIDVCENC HYPIVPLDGGK GTLLIRNGSE VRDPLVITYEG
61 SNPPASPLQD NLVIALHSYE PSHDGDGDFE KGEQLRILEQ SGewwkaQSL TTGQEGFIPF
121 NFVAKANSLE PEPWFFKNLS RKDAERQLLA PGNTHGSFLI RESESTAGSF SLSVRDFDQN
181 QGEVVKHYKI RNLDNGGFYI SPRITFPGLH ELVRHYTNAS DGLCTRLSRP CQTQKPQKPW
241 WEDEWEVPRE TLKLVERLGA GQFGEVWMGY YNGHTKVAVK SLKQGSMSPD AFLAEANLMK
301 QLQHQRLLVRL YAVVTQEPIY IITEYMENGS LVDFLKTSPG IKLTINKLLD MAAQIAEGMA
361 FIEERNYIHR DLRAANILVS DTLSCKIADF GLARLIEDNE YTAREGAKFP IKWTAPEAIN
421 YGTFTIKSDV WSFGILLTEI VTHGRIPYPG MTNPEVIQNL ERGYRMVRPD NCPEELYQLM
481 RLCWKERPED RPTFDYLRV LEDFFTATEG QYQPQP
```

Recombinant Lck nucleotide sequence:

```
1 atgcatcacc atcaccatca tatgggctgt ggctgcagct cacaccgga agatgactgg
61 atggaaaaca tcgatgtgtg tgagaactgc cattatccca tagtcccact ggatggcaag
121 ggcacgctgc tcatccgaaa tggctctgag gtgcgggacc cactggttac ctacgaaggc
181 tccaatccgc cggcttcccc actgcaagac aacctgggta tcgctctgca cagctatgag
241 ccctctcacg acggagatct gggctttgag aagggggaac agctccgcat cctggagcag
301 agcggcgagt ggtggaaggc gcagtccttg accacgggac aggaaggctt catccccttc
361 aatthttgtgg ccaaagcgaa cagtctggag cccgaaccct ggthcttcaa gaacctgagc
421 cgcaaggacg cggagcggca gctcctggcg cccgggaaca ctcacggctc cttcctcatc
481 cgggagagcg agagcaccgc gggatcgttt tcaactgtcgg tccgggactt cgaccagaac
541 cagggagagg tgggtgaaaca ttacaagatc cgtaactctgg acaacgggtg cttctacatc
601 tcccctcgaa tcacttttcc cggcctgcat gaactgggcc gccattacac caatgcttca
661 gatgggctgt gcacacgggt gagccgccc tgccagacc agaagcccca gaagccgtgg
721 tgggaggacg agtgggaggt tcccaggag acgctgaagc tgggggagc gctgggggct
781 ggacagttcg gggaggtgtg gatgggttac tacaacgggc acacgaaggt ggcggggaag
841 agcctgaagc agggcagcat gtccccggac gccttcctgg ccgaggccaa cctcatgaag
901 cagctgcaac accagcggct ggttcggctc tacgctgtgg tcaaccagga gccatctac
961 atcatcactg aatacatgga gaatgggagt ctagtggatt ttctcaagac cccttcaggc
1021 atcaagttga ccatcaaaa actcctggac atggcagccc aaattgcaga aggcattggca
1081 ttcattgaag agcgggaatta tattcatcgt gaccttcggg ctgccaacat tctgggtgtct
1141 gacaccctga gctgcaagat tgcagacttt ggcctagcac gcctcattga ggacaacgag
1201 tacacagcca gggagggggc caagtttccc ataatgtgga cagcgccaga agccattaac
1261 tacgggacat tcaccatcaa gtcagatgtg tggctctttg ggatcctgct gacggaaatt
1321 gtcacccacg gccgcatccc ttaccaggg atgaccaacc cggaggtgat tcagaacctg
1381 gacgagggct accgcatggt gcgcctgac aactgtccag aggagctgta ccaactcatg
1441 aggctgtgct ggaaggagcg cccagaggac cggcccacct ttgactacct gcgcagtgct
1501 ctggaggact tcttcacggc cacagagggc cagtaccagc ctcagccttg a
```

Reviewed and approved by site quality representative.

Unless otherwise stated in our catalogue or other company documentation accompanying the product(s), our products are intended for research use only and are not to be used for any other purpose, which includes but is not limited to, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses or any type of consumption or application to humans or animals.

©2007: Millipore Corporation. All rights reserved. No part of these works may be reproduced in any form without permission in writing.