



CHEMICON International, Inc. 28835 Single Oak Drive • Temecula, CA 92590
Phone: 909-676-8080 • 800-437-7500 • Fax: 909-676-9209
custserv@chemicon.com • techserv@chemicon.com • www.chemicon.com

Cy2 CONJUGATED AFFINITY PURIFIED SECONDARY ANTIBODY

- DESCRIPTION:** Whole molecule, absorbed for dual labeling, or F(ab')₂ fragment affinity purified antibody, Cy2 conjugated.
- QUANTITY:** Indicated on individual vials
- PURIFICATION:** This antibody was isolated from antisera by immunoaffinity chromatography using antigens coupled to agarose beads.
- FLUOROPHORE/
PROTEIN:** Approximately 1.9 (A492/A280)
- WAVELENGTH:** Absorption peak=492 nm, Emission peak= 510 nm
- APPLICATIONS:** Suggested dilution for most applications: 1:50-1:200
Optimal working dilutions must be determined by the end user.
- FORMAT:** Lyophilized. Buffer = 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6 with 15 mg/mL BSA, 0.01% Thimerosal and 0.05% sodium azide.
- RECONSTITUTION:** Reconstitute to 1 mg/mL with dH₂O
- STORAGE:** Maintain lyophilized product at 2-8°C for up to 12 months. After reconstitution the product is stable for several weeks at 2-8°C as an undiluted liquid. For extended storage after reconstitution, add an equal volume of glycerol to make a final concentration of 50% glycerol (ACS or better grade) followed by storage at -20°C in undiluted aliquots for up to 6 months. Please note the concentration of protein (and buffer salts) will decrease to one-half of the original after the addition of glycerol. Avoid repeated freeze/thaw cycles.
- WARNING:** For cyanine dye-labeled tissue or cells, use of mounting media containing phenylenediamine as an anti-fading reagent may result in weak or diffused fluorescence after storage of stained slides.

For research use only; not for use as a diagnostic.