

**GOAT ANTI-HUMAN IgG (Fc)
AFFINITY PURIFIED SECONDARY ANTIBODY
FITC CONJUGATE**

CATALOG NUMBER: AP113F

LOT NUMBER:

QUANTITY: 2 mg

CONCENTRATION: 2 mg/mL

SPECIFICITY: Shows a single band when run in an IEP system at a minimum concentration of 10 mg/mL, using human whole serum as the antigen. Reacts with Human IgG Fc fragment. No reactivity to other human immunoglobulins is observed.

**FLUOROPHORE/
PROTEIN:** 5.5 µg/mg

APPLICATIONS: Suggested dilution for most applications: 1:50-1:200
Optimal working dilutions must be determined by the end user.

FORMAT: Immunoglobulin purified by immunoaffinity chromatography conjugated to FITC.

PRESENTATION: Liquid 0.02M Phosphate buffer, 0.25M NaCl, pH 7.6 with 15 mg/mL BSA, 0.1% sodium azide.

STORAGE: Maintain at 2-8°C in undiluted aliquots for up to 6 months. DO NOT FREEZE. Store in the dark.

For research use only; not for use as a diagnostic.

© 2003-2005: CHEMICON International, Inc. - By CHEMICON International, Inc. All rights reserved. No part of these works may be reproduced in any form without permissions in writing.

USA & Canada • Phone: +1(800) 437-7500 • Fax: +1 (951) 676-9209 • Europe +44 (0) 23 8026 2233
Australia +61 3 9839 2000 • Germany +49-6192-207300 • ISO Registered worldwide
www.chemicon.com • custserv@chemicon.com • techserv@chemicon.com