

**MOUSE ANTI-NERVE GROWTH FACTOR-RECEPTOR, p75  
ALEXA FLUOR® 488 CONJUGATED  
MONOCLONAL ANTIBODY**

---

<b>CATALOG NUMBER:</b>	MAB5386X	<b>QUANTITY:</b>	50 µg
<b>LOT NUMBER:</b>		<b>CONCENTRATION:</b>	1.0 mg/mL
<b>ALTERNATE NAMES:</b>	low affinity NGF-Receptor; p75, NGFR	<b>EPITOPE:</b>	within aa 1-160, extracellular domain of human p75.
<b>CLONE NAME:</b>	ME20.4	<b>HOST/ISOTYPE:</b>	Ms IgG <sub>1</sub>
<b>BACKGROUND:</b>	The low affinity NGFR (Nerve growth factor receptor) is a 75kDa membrane-spanning glycoprotein lacking intrinsic tyrosine kinase activity. p75NGFR interacts with TrkA, the high affinity NGF receptor and potentiates TrkA signaling at low NGF concentrations. The p75 receptor binds nerve growth factor, brain-derived neurotrophic factor, neurotrophin-3 and neurotrophin-4 with varying specificities. The p75NGFR plays an important role in neurotrophic factor signaling and has been shown to modulate the susceptibility of selective cellular populations to programmed cell death. It is expressed on many neuronal cells types including many embryonic forms and the receptor can be used to isolate neuronal progenitor cells.		
<b>SPECIFICITY:</b>	Reacts with the low affinity nerve growth factor receptor (NGFR p75). The epitope recognized by the antibody maps within the third or fourth cysteine-rich repeats of the NGFR molecule. Reactivity has not been observed with rat, chicken, or mouse NGFR.		
<b>IMMUNOGEN:</b>	Human melanoma cell line (Ross, 1984).		
<b>APPLICATIONS:</b>	Immunocytochemistry Flow cytometry Optimal working dilutions must be determined by the end user.		
<b>SPECIES REACTIVITY:</b>	Reacts with Human, Rabbit, Porcine (Pig), and Sheep (Ovine). Does not react with Rat, Mouse and Chicken.		
<b>FORMAT:</b>	Purified immunoglobulin conjugated to Alexa Fluor® 488.		
<b>PRESENTATION:</b>	Liquid in PBS. Contains no preservative.		
<b>STORAGE/HANDLING:</b>	Aliquot and store at -20 °C, or glycerol (ASC grade or better) can be added (1:1) and the antibody stored at -20°C for up to 12 months from date of receipt. Avoid repeated freeze/thaw cycles.		
<b>REFERENCES:</b>	<ol style="list-style-type: none"><li>1. Ferreira, G, et al. (2001). Distribution and co-localization of choline acetyltransferase and p75 neurotrophin receptors in the sheep basal forebrain: implications for the use of a special cholinergic immunotoxin. <i>Neurosci.</i> 104:419-439.</li><li>2. Tremere, LA, et al. (2000). Antibody for human p75 LNTR identifies cholinergic basal forebrain of non-primate species. <i>Neuroreport</i> 11:2177-2183.</li></ol>		

---

3. Ross, AH, et al (1984). Characterization of nerve growth factor receptor in neural crest tumors using monoclonal antibodies. PNAS.USA 81:6681-6685.

**Important Note:** *During shipment, small volumes of product will occasionally become entrapped in the seal of the product vial. For products with volumes of 200  $\mu$ L or less, we recommend gently tapping the vial on a hard surface or briefly centrifuging the vial in a tabletop centrifuge to dislodge any liquid in the container's cap.*

***For research use only; not for use as a diagnostic.***

Unless otherwise stated in our catalog or other company documentation accompanying the product(s), our products are intended for research use only and are not to be used for any other purpose, which includes but is not limited to, unauthorized commercial uses, in vitro diagnostic uses, ex vivo or in vivo therapeutic uses or any type of consumption or application to humans or animals.

©2008: Millipore Corporation. All rights reserved. No part of these works may be reproduced in any form without permission in writing.

---

USA & Canada • Phone: +1(800) 437-7500 • Fax: +1 (951) 676-9209 • Australia +61 3 9839 2000  
www.millipore.com